$\Psi(2s)$ production in $p + A$ collisions: from SPS to FAIR

Partha Pratim Bhaduri
Variable Energy Cyclotron Centre, HBNI, 1/AF Bidhan Nagar, Kolkata 700 064, India

Abhijit Bhattacharyya
Department of Physics, University of Calcutta, 92, A. P. C. Road, Kolkata 700 009, India
(Dated: March 16, 2017)

We have analyzed the data on ψ' production in proton-nucleus ($p + A$) collisions, available from the NA50 Collaboration in the SPS energy domain. The investigated data sets include the absolute production cross sections as well as ψ'-to-Drell Yan (DY) cross section ratios. An adapted version of two component QVZ model has been employed to calculate ψ' production cross sections. For both ψ' and DY production, nuclear modifications to the free nucleon structure functions are taken into account. For ψ', final state interaction of the produced $c\bar{c}$ pairs with the nuclear medium is also taken into account, in accordance with the previously analyzed J/ψ data. A satisfactory description of the data in $p+A$ collisions is obtained. Model calculations are extrapolated to low energy regime to predict the ψ' suppression in proton induced collisions at FAIR SIS100.

arXiv:1703.04580v1 [nucl-th] 15 Mar 2017

I. INTRODUCTION

One of the focal aims of the relativistic heavy ion collision experiments is the quantitative mapping of the phase diagram of strongly interacting matter. Over past three decades enormous theoretical and experimental efforts have thus been devoted to study the properties of the hot and dense matter produced in such collisions. The major objective of the Compressed Baryonic Matter (CBM) experiment at FAIR is to explore the QCD phase diagram in the region of high net-baryon densities and moderate temperatures [1]. This experiment is designed to run at unprecedented high interaction rates which would enable precision measurements of rare diagnostic probes which are sensitive to the dense phase of the nuclear fireball. The FAIR Modularized Start Version (MSV) comprises the SIS100 ring which provides energies for gold beams up to 11 A GeV and for protons up to 30 GeV. The research program of CBM at SIS100 includes the systematic measurements of charmoniuam in $p + A$ collisions with varying target mass number, at proton energies up to 30 GeV. Such measurements are expected to shed light on the charmonium interaction with cold nuclear matter and constitute an important baseline for measurements in heavy ion collisions. Charmonium suppression in nuclear collisions is one of the oldest signatures to indicate the formation of quark-gluon plasma (QGP) [2]. However to identify the genuine plasma effects, a precise estimation of the suppression induced by the nuclear medium is a necessary prerequisite.

In our earlier work, we have predicted the J/ψ suppression in $p + A$ collisions at FAIR within the ambit of modified QVZ model. Different parameters of the model were fixed using the available data on J/ψ production in $p + p$ and $p + A$ collisions from different fixed target experiments. Model so calibrated was then extrapolated to the FAIR energy domain. In the present article, we plan to check the viability of our model by examining the available data corpus on ψ' production in $p + A$ collisions at SPS [3, 4]. After successful depiction of the SPS data, calculations are extrapolated to FAIR SIS100 energy regime.

II. A BRIEF DESCRIPTION OF THE MODEL

For calculation of ψ' production cross sections, we have used the adapted version of the originally proposed QVZ model [5]. Details of the original model and its modifications as used in the present calculations can be found in refs. [5–9]. Here we present a brief description for the sake of completeness. In hadronic collisions, ψ' production is assumed to be factorised into two steps. The initial stage is the perturbative production of the $c\bar{c}$ pair. This is followed by the non-perturbative formation of the physical resonance in the second stage. Resonance formation probabilities are incorporated in QVZ model using different functional forms which account for a wide variety of color neutralization mechanisms. The single differential ψ' production cross section in collisions of hadrons h_1 and h_2, at the center of mass energy $\sqrt{s}$ read as,

$$\frac{d\sigma^{\psi'}_{h_1h_2}}{dx_F} = K_{\psi'} \int dQ^2 \left(\frac{d\sigma^{c\bar{c}}_{h_1h_2}}{dQ^2 dx_F} \right) \times F_{c\bar{c}\to\psi'}(q^2), \quad (1)$$

where $Q^2 = q^2 + 4m_C^2$ with m_C being the mass of the charm quark and x_F is the Feynman scaling variable. $K_{\psi'}$ accounts for effective higher order contributions. $F_{c\bar{c}\to\psi'}(q^2)$ is the transition probability of a $c\bar{c}$ pair with relative momentum square q^2 to evolve into a physical ψ' meson. Two different functional forms namely Gaussian form ($F^{(\mathrm{G})}(q^2)$) and power law form ($F^{(\mathrm{P})}(q^2)$) mimicking two different mechanisms of color neutralization, have been found earlier to describe the absolute J/ψ production cross section data in $p + A$ collisions reasonably well [6]. Here we take the same two forms of $F(q^2)$, for calculating ψ' production cross section. They read as:



FIG. 1: (Color online) Model description of ψ' production in $p+A$ collisions at 400 and 450 GeV incident energy of the proton beams. Data are represented as the ratio of ψ' production cross sections in $p+A$ collisions to that in $p+Be$ collisions. As usual parctice, for 450 GeV beam energy, data for high intensity (HI) and and low intensity (LI) runs are separately shown. The two theoretical curves represents two different parametric forms of ψ' formation probability ($F(q^2)$).

$$F^{(\mathrm{G})}_{c\bar{c}\to\psi'(q^2)} = N_{\psi'}\,\theta(q^2)\,\exp\left[-q^2/(2\alpha_F^2)\right] \ , \qquad (2)$$

$$F^{(\mathrm{P})}_{c\bar{c}\to\psi'(q^2)} = N_{\psi'}\,\theta(q^2)\,\theta(4m_D^2-4m_C^2-q^2) \times \left(1-q^2/(4m_D^2-4m_C^2)\right)^{\alpha_F} \ , \qquad (3)$$

where m_D is the mass scale for the open charm production threshold. $N_{\psi'}$ and α_F are two parameters of $F_{c\bar{c}\to\psi'}(q^2)$, which should of course be different compared to those of J/ψ. They are fixed by comparing the model results with the available data. Introduction of $f_{\psi'}$ defined as $f_{\psi'} = K_{\psi'} \times N_{\psi'}$ helps in reducing the number of free parameters.

In $p+A$ collisions, ψ' production cross sections are modified due to cold nuclear matter effects. At the initial stage, nuclear modifications of the parton densities inside the target, modify the $c\bar{c}$ pair production cross section. In our analysis, we opted for leading order MSTW2008 [10] set for free proton pdf and EPS09 [11] interface to account for the nuclear effects. A detailed account of these initial state modifications of the parton distributions, as they are implemented in our model calculations can be found in ref. [6].

The nascent $c\bar{c}$ pairs interact with nuclear medium during their passage through the target. As a result the pairs gain momentum. In this process, some of the $c\bar{c}$ pairs can gain enough momentum to cross the threshold to transmute to open charm mesons. This results into the reduction in ψ' yield compared to the nucleon-nucleon collisions. The overall effect of the multiple scattering of the $c\bar{c}$ pairs inside the target is represented by a shift of q^2 in the transition probability,

$$q^2 \longrightarrow \bar{q}^2 = q^2 + \varepsilon^2 \ < L > \ . \qquad (4)$$

where ε^2 is the increase in the square of four momentum per unit path length. The mean geometrical path length $< L >$ is calculable within Glauber model.

TABLE I: Values of the fitting parameter α_F, for both Gaussian (G) and power-law (P) parametrizations, as fixed from ψ' production cross sections in $p+A$ collisions at SPS

Data set	$\alpha_F^{(G)}$	$\alpha_F^{(P)}$
NA50-400	$1.067 \pm 6.06 \times 10^{-2}$	$1.51 \pm 2.8 \times 10^{-1}$
NA50-450-HI	$1.09 \pm 7.22 \times 10^{-2}$	$1.41 \pm 3.12 \times 10^{-1}$
NA50-450-LI	$1.21 \pm 1.75 \times 10^{-1}$	$0.954 \pm 5.35 \times 10^{-1}$

Incorporation of final state dissociation in QVZ framework is largely different from the conventional approach. In usual practice, the nuclear dissociation of the different charmonium states are treated within Glauber model with an absorption cross section σ_{abs} quantifying the amount of dissociation. Note that, in the present approach, the nuclear effects are operative on the pre-resonant $c\bar{c}$ pairs which are yet to be hadronized. Hence for a given kinematic domain, the value of ϵ^2 should be same for both J/ψ and ψ' formation. We have seen in ref. [6] that for both parameterizations of $F(q^2)$, the corresponding values of ε^2 exhibited non-trivial dependence on the beam energy (E_b) [6]. Lower the beam energy, larger is the value of ε^2, implying larger nuclear dissociations.

III. ANALYSIS OF SPS DATA

With this brief description of the model, we now move forward to examine the viability of the model calculations in describing the available data on ψ' production in low energy $p+A$ collisions. Due to much smaller production cross sections, measurements are not as abundant as J/ψ



FIG. 2: (Color online) Model description of the data on ratio of the ψ'-to-Drell-Yan production cross-sections in 450 GeV and 400 GeV $p+A$ collisions, as published by NA50 Collaboration at SPS.



FIG. 3: (Color online) Model description of the data on the double ratio of the ψ'-to-J/ψ production cross-sections in 450 GeV and 400 GeV $p+A$ collisions. The double ratios are derived from the measured J/ψ and ψ' cross sections in $p+A$ collisions.

production, particularly at low energy fixed target experiments. At SPS a systematic measurement is available only from NA50 Collaboration, which measured ψ' production via their di-muon decay channel in 450 GeV [4] and 400 GeV [3] $p+A$ collisions. With 450 GeV proton beam, NA50 collected two sets of $p+A$ data in two independent runs with different beam intensities. The first set were collected with a high intensity (HI) 450 GeV proton beam using five different targets (Be, Al, Cu, Ag, W). Subsequently at the same beam energy new $p+A$ data samples were collected with same set of targets with low beam intensity (LI) and having 20 - 30% of the statistics of the HI set. Though they were initially collected in a slightly different kinematic domain, for a coherent comparison all the data sets at 450 GeV are corrected for a common phase space window: $-0.5 < y_{cms} < 0.5$ and $-0.5 < cos(\theta_{cs}) < 0.5$, where y_{cms} denotes di-muon centre-of-mass rapidity and θ_{cs} is the Collins-Soper angle. At 400 GeV, data were collected for six different targets (Be, Al, Cu, Ag, W, Pb) in the kinematic domain: $-0.425 < y_{cms} < 0.575$ and $-0.5 < cos(\theta_{cs}) < 0.5$. Even though the beam intensities were slightly higher than that of 450 GeV HI data samples, avalibility of the short beam time prevented the collection of sizeable statistics. In 400 GeV run, unlike previous measurements, data for all targets were collected in the same data taking period leading to significant reduction in the systematic errors. Later NA60 Collaboration studied charmonium production in $p+A$ collisions with proton beams of 400 and 158 GeV. Due to limitation in statistics their measurements were confined to J/ψ production only.

Here a few words on the limitations of the analysis of ψ' production in low energy collisions may be in order. For describing the production in $p+A$ collisions, the

QVZ model has effectively three parameters namely $f_{\psi'}$ and α_F to account for hadronization and ϵ^2 to account for final state dissociation. Amongst them ϵ^2 is already fixed from our previous analysis of J/ψ production cross sections, as argued above. The other two parameters namely α_F and $f_{\psi'}$ are to be fixed from the present data. This is in contrast to our earlier J/ψ analysis where both $f_{J/\psi}$ and α_F were fixed from the analysis of inclusive J/ψ production cross sections as a function of beam energy in $p+p$ collision. Unfortunately no such data is available for ψ' production, leaving us to determine all the free parameters from $p+A$ data only.

Fig. 1 shows the variation of the ψ' production cross section for different target nuclei, as a function of $< L >$, in $p+A$ collisions measured by NA50 Collaboration. The data are expressed as the ratio of the ψ' production cross sections in $p+A$ collisions to that in $p+Be$ collisions. One advantage of fitting this ratio is that we get rid of the multiplicative fitting parameter $f_{\psi'}$. The two theory curves result from fitting the above data sets following two parameterisations of $F(q^2)$. The $< L >$ values, as published with the data, are used to generate the theoretical curves. The α_F values extracted from different data sets are given in Table I. They are comparable within errors *i.e.* they show a very weak beam energy dependence. We have also fitted the ψ'-to-Drell-Yan ratio as shown in Fig. 2. In Drell-Yan (DY) process, a quark and an antiquark from the nucleons of the two colliding nuclei annihilate to form a virtual photon which subsequently decays into a $\mu^+\mu^-$ pair. We have calculated the leading order DY cross section using the standard prescription [12]. The only CNM effect incorporated is the nuclear modification of the quark distributions inside the target. The inclusive cross sections are obtained by integrating the double differential cross section within the suitable mass and rapidity range as appropriate for a particular data set. Like our previous analysis, the only free parameter in this case is K_{eff} defined as $K_{eff} = f_{\psi'}/K_{DY}$ where $f_{\psi'} = K_{\psi'} \times N_{\psi'}$. K_{DY} takes care of higher order effects in DY production. The α_F values are same as obtained from the fitting of the ratio $\sigma^{\psi'}_{p+A}$ to $\sigma^{\psi'}_{p+Be}$.

Finally we describe the double ratio for ψ'-to-J/ψ production cross sections in Fig. 3. No parameter is tuned for this observable. The experimental data points are deduced from the measured values accounting for the appropriate propagation of errors. The theoretical curves result from the model analysis of the single ratios ($p+A$ to $p+Be$) for J/ψ and ψ'. As expected both the curves give a satisfactory description of the data.

IV. PREDICTION AT FAIR

The basic motivation of this work is to understand ψ' production at SPS and then to extrapolate it to a much denser scenario which is relevant for nuclear collisions at FAIR. In future, FAIR is going to provide us a plethora of data at high densities. At such high densities, a clear understanding of charmonium production is expected to shed light on the phase diagram of strongly interacting matter. With that motivation we now extrapolate our model calculations to estimate the ψ' production expected in $p+A$ collisions at FAIR SIS100 synchrotron. To do this we need to fix the values of α_F at relevant beam energies. As we saw in the previous section the α_F values obtained from fitting the SPS data are comparable within error bars. We have fitted those values to deduce beam energy independent constant values $\alpha_F^{(G)} = 1.085 \pm 4.49 \times 10^{-2}$ and $\alpha_F^{(P)} = 1.4 \pm 1.94 \times 10^{-1}$. These α_Fs are then used to estimate the cross sections at FAIR. The beam kinematic threshold for ψ' production in elementary collisions is $E_b^{Th} \simeq 15.6$ GeV. Thus ψ' production may only be significant at top SIS100 energy. Fig 4 represents the ψ' production cross section in 30 GeV $p+A$ collisions for different nuclear targets. Result is expressed in terms of the ratio of inclusive production cross sections, evaluated over a rapidity slice $-0.5 \leq y_{cms} \leq 0.5$. Beam energy dependence of $\epsilon^2(E_b)$, as parametrized in [6] is used to estimate the final state dissociation. The amount of suppression is larger than that measured at SPS. As investigated in ref. [13], this is caused by convolution of two CNM effects the shadowing of the parton densities inside the target nuclei and the amplification of final state dissociation in the kinematic region probed at FAIR. Note that in our model both the CNM effects are same for both J/ψ and ψ'. In the spirit of Glauber analysis, we have fitted the suppression curves with exponential parametrization with an effective absorption cross section σ_{abs} as $\sigma_{pA} = \sigma_0 e^{-\rho_0 \sigma_{abs} < L >}$. This leads to $\sigma_{abs} \simeq 12$ mb for Gaussian and $\sigma_{abs} \simeq 11$ mb for power-law parametrizations. At SPS, suppression data in $p+A$ collisions were also expressed in terms of ψ'-to-DY ratio. Being a purely electromagnetic process, the cross section for DY production would be orders of magnitude smaller than that of charmonia and detection of DY di-muons may not be feasible at FAIR.



FIG. 4: (Color online) Model description of ψ' production in 30 GeV $p+A$ collisions at FAIR. Two theoretical curves represent two different mechanisms of resonance formation.

Before we close, limitations on the applicability of our

model calculations at FAIR energy domain may be noted. QVZ model is based on QCD factorization which in general is operative at high energies and we assume that factorization still holds for charmonium production close to threshold. An alternative approach based on scaling arguments for near threshold quarkonium production can be found in [14]. Moreover in the model the resonance formation is believed to occur outside the nuclear medium. If the resonances are formed inside the nuclear matter, their final state dissociation might be different. Charmonium data at FAIR SIS100 are thus highly welcome to validate or nullify some of the above arguments.

V. SUMMARY

In summary, we have analyzed the available SPS data on ψ' production in $p + A$ collisions measured by NA50 Collaboration. The adapted version of QVZ model is employed for this purpose. The model has been found earlier to give reasonable description of J/ψ production at SPS. Non-availability of suitable data makes the analysis rather complex by adding large uncertainites in determination of the model parameters. For both ψ' and DY initial state modification of the parton distributions inside the target are taken into account. Within the QVZ approach, the final state dissociation of the ψ' mesons inside the cold nuclear medium is accounted through the multiple scattering of the pre-resonant $c\bar{c}$ pairs with the spectator nucleons. The model is found to describe the observed suppression for both forms of the transition probability. Model calculations are extrapolated to the FAIR energy domain. A much larger suppression is foreseen at SIS100 $p + A$ collisions. The preliminary runs at FAIR should be able to remove large uncertainties in the parameters and thus can lead to a much precise estimate. With such precise estimates in $p + A$ scenario one can plan to extrapolate to $A+A$ scenario once SIS300 synchrotron ring becomes operative.

Acknowledgement

AB would like to thank UGC(DRS) and BRNS for support.

[1] T. Ablyazimov *et al.* CBM Collaboration, arXiv:1607.01487 [nucl-ex].
[2] T. Matsui and H. Satz, Phys. Lett. B**178**, 416 (1986).
[3] B. Alessandro *et al.* NA50 Collaboration, Euro. J.Phys **48** 329 (2006).
[4] B. Alessandro *et al.* NA50 Collaboration, Euro. J.Phys **33** 31 (2004).
[5] J. Qui, J.P. Vary and X. Zhang, Nucl. Phys. A698, 571 (2002); Phys. Rev. Lett. **88** 232301 (2002).
[6] P. P. Bhaduri, A.K. Chaudhuri and S. Chattopadhyay, Phys. Rev. C **84**, 054914 (2011).
[7] P. P. Bhaduri, A. K. Chaudhuri and S. Chattopadhyay, Phys. Rev.C **85**,064911 (2012).
[8] A. K. Chaudhuri, Phys. Rev. Lett. **88**, 232302 (2002).
[9] A. K. Chaudhuri, Phys. Rev.C **66**,021902 (2002).
[10] A.D. Martin,W.J. Stirling, R.S. Thorne, G. Watt, Eur.Phys.J.C63:189-285,2009; A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, Eur.Phys.J.C64:653-680,2009; A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, Eur.Phys.J.C70:51-72,2010.
[11] K.J. Eskola, H. Paukkunen and C.A. Salgado, JHEP04 (2009) 065.
[12] R. Vogt, Physics Reports 310, 197 (1999);
[13] P. P. Bhaduri, A. K. Chaudhuri and S. Chattopadhyay, Phys. Rev.C **89**,044912 (2014).
[14] P. P. Bhaduri and S. Gupta, Phys. Rev.C **88**,045205 (2013).